

The synaptic muscle-specific kinase (MuSK) complex: new partners, new functions

Laure Stochlic, Annie Cartaud, and Jean Cartaud*

Summary

The muscle-specific kinase MuSK is part of an agrin receptor complex that stimulates tyrosine phosphorylation and drives clustering of acetylcholine receptors (AChRs) in the postsynaptic membrane at the vertebrate neuromuscular junction (NMJ). MuSK also regulates synaptic gene transcription in subsynaptic nuclei. Over the past few years, decisive progress has been made in the identification of MuSK effectors, helping to understand its function in the formation of the NMJ. Similarly to AChR, MuSK and several of its partners are the target of mutations responsible for diseases of the NMJ, such as congenital myasthenic syndromes. This minireview will focus on the multiple MuSK effectors so far identified that place MuSK at the center of a multifunctional signaling complex involved in the organization of the NMJ and associated disorders. *BioEssays* 27:1129–1135, 2005. © 2005 Wiley Periodicals, Inc.

Introduction

The function of chemical synapses is based on the accumulation of ligand-gated channels or receptors for neurotrans-

mitters in the postsynaptic membrane domain. At the neuromuscular junction (NMJ), nicotinic AChR accumulates in the crests of the postsynaptic membrane with a surface density of 10–20,000/ μm^2 directly beneath the nerve terminal.^(reviewed in 1) Extracellular, transmembrane and cytoplasmic proteins, which have structural, adhesive and signaling roles, coaggregate with AChRs in the postsynaptic specialization. The synaptic specialization extends to the basal lamina, which is locally enriched in specific isoforms of laminin, in perlecan and in acetylcholinesterase (AChE, EC 3.1.1.7).^(reviewed in 2) The synaptic basal lamina also contains two major factors expressed and secreted by motor nerves, agrin⁽³⁾ and neuregulins (NRG),^(reviewed in 4) which are both implicated in synapse formation and maintenance.^(reviewed in 1,5) NRG-1 activates ErbB receptors (Receptor Tyrosine Kinase belonging to the EGF receptor family) in the postsynaptic membrane and is thought to trigger transcription of AChR and other synaptic genes. Agrin is a large heparansulfate proteoglycan, first isolated by McMahan and colleagues from the Torpedo electric organ, on the basis of AChR aggregating activity in vitro.⁽³⁾ Several agrin isoforms that differ in their capacity to induce AChR clustering are generated by alternative splicing, the most-active forms being expressed by motoneurons during embryogenesis precisely during the innervation phase of muscle development.^(reviewed in 6) Most interestingly, ectopic postsynaptic specializations are induced in innervated myofibers following injection and expression of plasmids coding for neural agrin.⁽⁷⁾ The other important component for the differentiation of the NMJ is MuSK, a receptor tyrosine kinase specific for muscle cells,^(8,9) expressed and concentrated in the postsynaptic membrane in Torpedo electrocyte and in skeletal muscle.^(8–10) Mice deficient for MuSK or agrin have similar phenotypes and lack normal neuromuscular synapses at birth.^(11,12) These results together with data from previous experiments, which show that MuSK is activated by agrin, suggest that MuSK is a component of an agrin receptor complex. Agrin was initially believed to promote the clustering of AChRs at the NMJ without major effects on transcriptional regulation. However, several studies suggest that agrin could also stimulate the transcription of synaptic genes and that MuSK is required for synapse-specific transcription.^(7,13–20) MuSK-deficient mice also disclose presynaptic defects in addition to postsynaptic ones, indicating that MuSK activation

Biologie Cellulaire des Membranes, Institut Jacques Monod, Paris, France

Funding agency: This work was supported by the Centre National de la Recherche Scientifique, the Universities Paris VI & Paris VII, and the Association Française contre les Myopathies (AFM).

*Correspondence to: Dr. Jean Cartaud, Biologie Cellulaire des Membranes, Institut Jacques Monod, UMR 7592, CNRS/Universités Paris VI et VII, 2 Place Jussieu, 75251 Paris Cedex 05, France.

E-mail: cartaud@ijm.jussieu.fr

DOI 10.1002/bies.20305

Published online in Wiley InterScience (www.interscience.wiley.com).

Abbreviations: Abl, Abelson kinases; AChR, acetylcholine receptor; AChE, acetylcholinesterase; APC, Adenomatous Polyposis Coli; ColQ, collagenic tail of AChE; Dvl, Dishevelled; GABP, GA binding-protein; GGT, geranylgeranyltransferase; JNK, c-jun NH₂-terminal kinase; MAGI, membrane-associated guanylate kinase (MAGUK) with inverted domain orientation; MAPK, mitogen-activated protein kinase; MuSK, muscle-specific kinase; NMJ, neuromuscular junction; NRG, neuregulin-1; PDZ, Postsynaptic density, Disc-large and Zonula occludens. HUGO gene symbols of human genes involved in NMJ disorders are: acetylcholinesterase, AChE; acetylcholine receptor (α , β , γ , δ , ε subunits), CHRN(A-E); ColQ, COLQ; MuSK, MUSK.


is also required for presynaptic differentiation. This process involves retrograde signals so far unidentified.

Despite many efforts, the direct physical interaction between MuSK and agrin has so far not been clearly established and the mechanism of activation of MuSK remains obscure. An associated component selectively expressed in skeletal muscle cells, termed MASC (Muscle-Associated Specific Component), has been postulated as coreceptor for agrin to activate MuSK.⁽²¹⁾ MASC has yet to be identified. Coligands for agrin or post-translational modifications (glycosylation) either of agrin or of the ectodomain of MuSK could also be envisaged as cofactors for agrin activation of MuSK.

The pleiotropic role of MuSK in synaptic differentiation likely relies on its interaction, either direct or indirect, with a wealth of downstream effectors. In the last few years, several partners of MuSK have been reported, shedding some light onto the mechanisms of action of MuSK. An integrated view of these various mechanisms, taking into account the most-recent discoveries in the field, is however still pending. In this review, we summarized recent advances in the function of the MuSK complex as a multifunctional platform regulating synapse formation and/or maintenance.

The MuSK signaling pathway involves rapsyn and tyrosine phosphorylations

Agrin stimulates the dimerization⁽²²⁾ and the kinase activity of MuSK, which is essential for clustering of AChRs.⁽²³⁾ Signaling triggered from MuSK activation depends upon the phosphorylation of a tyrosine residue (Y553) in the juxtamembrane consensus recognition site (NPXY) for phosphotyrosine-binding domain-containing proteins^(18,24) (Fig. 1). This phos-



phorytation is required to recruit downstream signaling components involved in many aspects of MuSK signaling, including tyrosine phosphorylation and AChR clustering as well as synapse-specific transcription.⁽¹⁸⁾ Furthermore, agrin causes rapid activation of Src-related kinases and tyrosine phosphorylation of AChR β and δ subunits,⁽²⁵⁾ the phosphorylation of β being required for efficient AChR clustering.⁽²⁶⁾ Tyrosine phosphorylation as well as Src and Fyn kinases are necessary for cluster stabilization.⁽²⁷⁾ However, MuSK kinase domain activation and the resulting phosphorylation of AChR subunits are not sufficient for AChR clustering, indicating that AChR phosphorylation is not the sole trigger of the clustering process.⁽²³⁾

Rapsyn (Receptor-associated protein at the synapse), the 43-kDa cytoplasmic peripheral membrane protein associated with AChRs in the postsynaptic membrane^(reviewed in 5) is an essential player in AChR aggregation. Rapsyn-deficient mice die at birth and, like MuSK mutants, lack all known features of postsynaptic differentiation, in particular AChR aggregates.⁽²⁸⁾ In these mutants, MuSK normally accumulates at the central region of muscle fibers, indicating that rapsyn is a required intermediate in the pathway that couples MuSK activation to AChR clustering, possibly by recruiting synaptic components to a MuSK-containing scaffold.⁽²⁹⁾ Unlike the ectodomain of all other receptor tyrosine kinases simply mediating ligand binding, several reports point to an additional role of the ectodomain of MuSK in AChR clustering.^(18,24,28) Mapping domains on MuSK necessary for its interaction with rapsyn indicate that sequences in and near the fourth IgG-like domain (see Fig. 1) are required for association with rapsyn through a putative protein RATL (Rapsyn Associated Transmembrane Linker) to mediate AChR aggregation in quail QT-6 cells.^(24,29) Examination of agrin-inducing signaling events in rapsyn-deficient myotubes reveals that the agrin-induced MuSK activation occurs normally, whereas MuSK-dependent phosphorylation of AChR (and AChR clustering) does not occur. AChR phosphorylation is thus rapsyn-dependent. Src kinase, the major kinase phosphorylating AChR has been shown to be associated with rapsyn.^(30,31) Rapsyn may thus function as a scaffolding protein presenting a MuSK-activated Src-related kinase to the AChRs.

MuSK triggers the clustering of synaptic proteins via actin cytoskeleton reorganization

Convergent studies highlight the participation of actin in the process of AChR aggregation following agrin activation of MuSK. Agrin stimulation results in actin polymerization and translocation of the actin regulatory protein cortactin to postsynaptic sites.⁽³²⁾ Moreover, latrunculin A, an inhibitor of actin polymerization blocks agrin-induced AChR clustering in myotubes.⁽³²⁾ The Rho GTP-ases Rac/Cdc42, two ubiquitous monomeric small GTPases regulating actin organization, are required for agrin-elicited clustering of AChRs in myotubes.⁽³³⁾

Furthermore, Rac and Rho GTP-ases play distinct but complementary roles in the mechanism of AChR clustering. Rac is necessary for the initial phase of formation of microclusters, and Rho required for their subsequent condensation into full-size AChR clusters.⁽³⁴⁾


The recent discovery of several MuSK effectors has shed some light on the molecular events downstream of agrin-induced MuSK activation (Fig. 2). Interestingly, several of these effectors are involved in actin cytoskeleton organization. In addition to the MuSK kinase activity, a nonreceptor tyrosine kinase (NRTK) activity placed downstream of MuSK is necessary for AChR clustering. Pendergast and collaborators⁽³⁵⁾ hypothesized a role for Abelson tyrosine kinases (Abl) in synapse formation and reported the requirement of these proteins in agrin-induced AChR clustering in vitro. Abl1 and Abl2 (also known as Abelson and Arg respectively) define a family of NRTK characterized by several domains, including SH2 and SH3, a tyrosine kinase and a unique C-terminal actin-binding site. Abl kinases regulate actin organization and

presynaptic axon guidance.⁽³⁶⁾ Abl kinases are localized to the postsynaptic membrane of the developing NMJ. MuSK and Abl kinases effected reciprocal tyrosine phosphorylation and formed a complex upon agrin engagement. However, the binding site for Abl kinase on the cytoplasmic domain of MuSK remains to be determined. The authors proposed that Abl kinases amplify initial signaling and AChR clustering through tyrosine phosphorylations and stabilize clusters via induction of synaptic actin network.⁽³⁵⁾

Using the yeast two-hybrid system with the cytoplasmic region of MuSK as a bait, Mei and collaborators⁽³⁷⁾ identified Dishevelled 1 (Dvl), a scaffolding protein originally discovered in *Drosophila* that is implicated in the development of coherent arrays of polarized cells via the Wnt signaling pathway. Dvls are cytoplasmic proteins comprising three domains: DIX (Dishevelled-Axin), PDZ (Postsynaptic density/Disc large/ZO-1) and DEP (Dishevelled-Egl-10-Pleckstrin). The authors demonstrated that Dvl binds to both the juxtamembranous and kinase domains of MuSK through its DEP domain. However, this association is independent of MuSK phosphorylation and the disruption of the MuSK–Dvl interaction has a modest effect (= two-fold decrease) on agrin-induced AChR clustering. Dvl also interacts with the serine/threonine kinase PAK1 (p21-Activated Kinase), which becomes activated in myotubes in response to agrin. PAK1 is an effector of Rac and Cdc42 in actin reorganization. Dvl, by forming a scaffold with MuSK and PAK1, would thus regulate AChR clustering mediated by agrin.⁽³⁷⁾ Yet, the confirmation of the role of Dvl in the formation of the NMJ in vivo is awaited.

Another potentially important signaling component in the agrin/MuSK pathway is geranylgeranyltransferase 1 (GGT, EC 2.5.1.10). Double hybrid experiments shown that the α subunit of GGT interacts with the kinase domain of MuSK.⁽³⁸⁾ Agrin causes rapid increase in tyrosine phosphorylation of GGT, and inhibition of either GGT expression or of GGT activity prevents muscle cells from forming AChR clusters in response to agrin. GGT inhibitors prevent agrin-induced activation of Rac1, Cdc42 and PAK1 in myotubes. Moreover, transgenic mice expressing an inactive GGT mutant still able to bind MuSK, exhibit NMJ defects. Strictly, these data indicate that prenylation (a post-translational modification of proteins catalyzed by GGT conferring membrane targeting) of key components of the MuSK signaling pathway, for example Rac, is necessary for AChR clustering. Since the MuSK–GGT association is independent of MuSK phosphorylation, a direct demonstration of the role of MuSK–GGT interaction is still pending.

Evidence for a regulation of AChR clustering by the tumor suppressor APC (Adenomatous Polyposis Coli), an actin-binding protein, has been reported.⁽³⁹⁾ The APC protein has been found in several subcellular compartments of mammalian and *Drosophila* cells, including the cytoplasm, nucleus and adhesive junctions. APC colocalizes with AChRs in the postsynaptic membrane of the NMJ and directly interacts with



the β subunit of AChRs, helping to localize them to the actin cytoskeleton. Another actin-binding complex, the utrophin glycoprotein complex, is specifically enriched in the postsynaptic membrane.^(reviewed in 40) Although not directly involved in the early steps of the AChR clustering process, this complex could play a role in the stabilization of AChR clusters, possibly linking rapsyn to the actin cytoskeleton through dystroglycan.⁽⁴¹⁾ One component of the utrophin complex, the phosphoprotein dystrobrevin,⁽⁴²⁾ a substrate for tyrosine kinases, is implicated in the stability of agrin-induced AChR clusters.⁽⁴³⁾

Finally, the scaffolding protein, MAGI-1c, has been identified by a proteomic approach as a MuSK partner in the postsynaptic membrane in Torpedo electrocytes.⁽⁴⁴⁾ MAGI-1c⁽⁴⁵⁾ is a member of the MAGUK family of scaffolding proteins involved in cell polarity and organization of signal transduction within cellular junctions, excitatory central synapses and the NMJ in *Drosophila*.^(reviewed in 46) As for other MAGUKs associated with synapses, e.g. the PSD-95/SAP90 and S-SCAM, MAGI-1c is composed of several protein–protein interaction domains: a guanylate kinase (GUK) domain, two WW domains and six PDZ domains. Each one of these domains potentially confers specific association with a large panel of signaling, structural and cytoskeletal elements. MAGI-1c is localized at the Torpedo electrocyte cholinergic synapses and at the adult rat NMJs but was not detected in agrin-elicited AChR clusters in myotubes, suggesting that MAGI-1c is not involved in the clustering mechanism. In line with this observation, pull-down experiments revealed that the PDZ 4/5 domains of MAGI-1c interact with the consensus C-terminal PDZ-binding site of MuSK (Cartaud et al., unpublished observations), a site that is dispensable for its clustering activity in cultured muscle cells.⁽²⁴⁾ Similarly, PSD-95 family proteins are not required for receptor clustering but are believed to be organizers of excitatory synapses, and are involved in synaptic plasticity.^(reviewed in 47) Likewise, MAGI-1c might be involved in the building up of a specialized protein complex at the NMJ participating to the terminal differentiation of the NMJ.

MuSK and the regulation of synaptic gene transcription

Genes encoding AChR subunits, as well as utrophin, rapsyn, AChE and MuSK are selectively transcribed in myofiber nuclei localized beneath the nerve contact. Thus, synapse-specific transcription is believed to participate in concentrating gene products at the NMJ.^(reviewed in 1,48) NRG⁽⁴⁾ acting through ErbB tyrosine kinase receptors in the postsynaptic membrane activates synaptic gene transcription via the Mitogen-Activated Protein Kinase (MAPK), c-jun NH₂-terminal kinase (JNK) and the Phosphatidylinositol-3 Kinase (PI3K) signal transduction pathways (see Fig. 3). These pathways converge to the phosphorylation of the Ets transcription factor GA binding-protein (GABP). Phosphorylated GABP and




Figure 3. Model for transcriptional regulation of synaptic genes at the NMJ. The tyrosine kinase receptors ErbB 2/4 and MuSK trigger several signaling cascades that cooperate to initiate transcriptional activation of synaptic genes in subsynaptic nuclei. Activation of ErbB receptors by neuregulin (NRG) triggers both the MAPK and JNK pathways via Ras. This leads to the expression of c-Jun and to the phosphorylation of the Ets-related transcription factor GABP that will activate transcription of synapse-specific genes through binding to the N-box (N). MuSK also induces activation of JNK via Rac/Cdc42, suggesting that the two pathways are connected. Inhibitory factors such as Erbin and SHP2, as well as 14-3-3 γ interacting with MuSK, inactivate the ErbB signaling pathway. The question mark indicates a possible regulation of the MuSK–14-3-3 γ interaction by a still unknown signal either acting on MuSK or on another signaling molecule. The PI3K pathway is not indicated in the model. (Modified from refs. 15,48).

JNK/c-jun activate transcription of the synaptic genes via GABP binding to the N-box, a 6-bp promoter segment CCGAA that directs synapse-specific gene expression.^(reviewed in 48)

Two negative regulators of the NRG/ErbB pathway, the protein tyrosine phosphatase SHP2 and erbin, have been identified. SHP2 dephosphorylates and inactivates ErbB receptors in the postsynaptic membrane after NRG stimulation and attenuates the NRG-induced expression of the *cAChR* promoter reporter gene in cultured myotubes.⁽⁴⁹⁾ Erbin associates with the C terminus of ErbB2 through its PDZ domain and downregulates the MAPK signaling cascade stimulated by NRG, through inactivation of Ras.⁽⁵⁰⁾

In addition to the NRG/ErbB pathway, several studies pointed to a role for agrin and MuSK in the regulation of synaptic gene transcription. Neural agrin or constitutively active MuSK cluster the ErbB receptors and stimulate transcription of AChR genes.^(7,51) Recently, Brenner and collaborators demonstrated that agrin-induced *musk* transcription is controlled in

part by the NRG/ErbB pathway organized by agrin–MuSK and by a novel shunt path in which MuSK stimulates the activation of Rac and JNK in a manner independent of the NRG/ErbB pathway. Both pathways converge onto the N-box in the *musk* and possibly other synaptic gene promoters.⁽¹⁵⁾ In fetal muscles, formation of postsynaptic-like sites and targeted gene expression begins to develop in the absence of agrin but depends on MuSK.^(19,20) Given these data, MuSK appears to be involved not only in AChR clustering but also in the regulation of synaptic genes expression. Interestingly, Rac/Cdc42 appear to play a central role in agrin/MuSK signaling conducting both to synaptic gene transcription and AChR clustering (see Figs. 2 and 3).

In a systematic search for MuSK partners in situ, Cartaud and collaborators⁽⁵²⁾ identified the adaptor protein 14-3-3 γ , using a proteomic approach in purified postsynaptic membrane from Torpedo electrocytes. The 14-3-3 γ protein colocalizes with AChRs at the NMJ in rat muscles and coimmunoprecipitates with MuSK. 14-3-3 proteins constitute an emerging family of signaling molecules present in eukaryotic organisms involved in intracellular signaling.^(reviewed in 53) Among many functions in cell signaling, 14-3-3 proteins are major regulators of the MAPK and PI3K signaling pathways. Overexpression of 14-3-3 γ in muscle cells specifically represses transcription of several synaptic genes including α , δ and ϵ -AChR subunits, *utrophin* (promoter A), *rapsyn*, and *MuSK*. The 14-3-3 proteins are known to interact and modulate the activity of Raf-1.⁽⁵⁴⁾ In myotubes in culture, 14-3-3 γ inhibits the activation of ϵ -AChR induced by constitutively active Raf-1, suggesting that it acts downstream of the NRG/ErbB pathway by blocking Raf-1 in an inactive conformation. Moreover, the expression of 14-3-3 γ in muscle fibers in vivo causes both the repression of synaptic genes transcription and morphological perturbations of the NMJ. Taken together, these data provide evidence that MuSK may serve as a scaffold to localize and concentrate 14-3-3 γ close to the postsynaptic membrane. At this location, 14-3-3 γ is prone to interact with signaling molecules of the NRG/ErbB cascade such as Raf-1 (see Fig. 3). In many cases, the interaction of 14-3-3 proteins with their various partners is regulated by phosphorylation of serine residues in consensus sequences in the target proteins.^(reviewed in 55) Two such sequences have been identified in the cytoplasmic region of rat MuSK.⁽⁵²⁾ An interesting hypothesis raised by these observations is that a signaling molecule, probably different from agrin, would regulate the MuSK–14-3-3 interaction, thereby regulating the synaptic transcription.

MuSK localizes asymmetric forms of acetylcholinesterase at the NMJ

At the NMJ, AChE is mainly present as asymmetric forms in which three tetramers of catalytic subunits are associated with a structural collagenic tail named ColQ.^(reviewed in 2) The

accumulation of the enzyme in the synaptic basal lamina strictly relies on ColQ. This mechanism is mediated by interaction between the collagenous domain of ColQ and perlecan, an heparan sulfate proteoglycan, which itself binds α -dystroglycan, a transmembrane glycoprotein of the dystrophin–utrophin complex enriched at the NMJ. In the absence of α -dystroglycan,⁽⁵⁶⁾ perlecan and AChE do not colocalize with AChR. Also, in perlecan null mice, AChE is absent from the NMJ.⁽⁵⁷⁾ These data suggest the requirement for the dystrophin–utrophin complex in clustering AChE at synaptic sites. However, recent studies indicate that the C terminus of ColQ is required for AChE anchoring at the NMJ,⁽⁵⁸⁾ suggesting that another protein might interact with AChE and contribute the specificity needed for synaptic localization. This protein has recently been identified as MuSK.⁽⁵⁹⁾ MuSK and ColQ can be crosslinked in situ in Torpedo AChR-rich membranes. Both proteins colocalize and coimmunoprecipitate in COS-7 cells cotransfected with plasmids encoding MuSK and ColQ. Moreover, MuSK is required to form cell surface AChE clusters in cultured myotubes.⁽⁵⁹⁾ Taken together, these observations suggest that a ternary complex containing ColQ, perlecan and MuSK is required for AChE clustering and demonstrate that MuSK dictates AChE synaptic localization at the NMJ. They also provide a rational molecular basis explaining certain forms of congenital myasthenic syndrome (CMS) associated with mutations in the C-terminal part of ColQ. Surprisingly, in these patients, the asymmetric forms of AChE are normally synthesized, assembled and secreted. Yet, no AChE accumulate in the synaptic cleft.^(reviewed in 60) The impossibility for these mutants of ColQ to bind efficiently with MuSK at synaptic sites is undoubtedly causal in these disorders.

MuSK: a new target for autoimmune and genetic disorders at the neuromuscular junction

Because of its importance in NMJ development and function, and similar to AChR and other synaptic proteins, MuSK is a possible target for autoimmune and genetic disorders at the NMJ.^(61,62) Myasthenia gravis is an antibody-mediated autoimmune disease of the NMJ caused, in the majority of patients, by the presence of auto-antibodies to AChR and the consecutive loss of AChRs in the postsynaptic membrane. In 70% of myasthenia gravis patients seronegative for AChR-antibodies, circulating auto-antibodies to MuSK are present.⁽⁶³⁾ These observations define immunologically distinct forms of the disease. The MuSK auto-antibodies were specific for the extracellular domains of MuSK and strongly inhibited MuSK function.⁽⁶³⁾ These antibodies are thus likely involved in the pathogenesis. Other major disorders of the NMJ—the congenital myasthenic syndromes – caused by mutations in key components of the postsynaptic membrane such as AChR subunits and AChE (see above) have been

reported.^(reviewed in 60) Recently, MuSK has been shown to be the direct target for mutations causing CMS associated with dramatic presynaptic and postsynaptic structural abnormalities and severe decreases in AChR ϵ -subunit and MuSK expression.⁽⁶²⁾ Mutations in rapsyn, a downstream effector of MuSK, also caused myasthenic syndromes.^(reviewed in 60) One could predict that novel mutations in MuSK will be discovered in myasthenic patients, and that MuSK effectors are potential molecular targets for neurological diseases at the NMJ. Understanding the fundamental molecular and cellular bases underlying MuSK function holds the promise of therapeutic approaches for treating these pathologies.

Conclusion

The increasing knowledge of the MuSK effectors provides a coherent molecular basis to the functional complexity of MuSK, sustaining its pleiotropic function in NMJ formation. In the model proposed (Fig. 2), upon stimulation by agrin, MuSK recruits Abl kinases which activate Rac/Cdc42 which would in turn stimulate PAK1 and other targets of activated GTPases. Dvl, owing to its interaction with MuSK and PAK1, functions as a scaffold molecule bridging the two proteins to form a signaling complex that leads to local actin reorganization, ultimately leading to AChR clustering. In this context, an important role for GGT would be to confer membrane localization of Rac/Cdc42, and thus basal Rac/Cdc42 activation. However, the steps in agrin signaling between MuSK activation and actin cytoskeleton anchoring of AChRs via rapsyn are still uncompletely characterized.

Other key MuSK partners, 14-3-3 γ and ColQ, fulfil different dedicated roles in synapse formation and/or maintenance, such as the regulation of synaptic gene expression and the localization of AChE, respectively. How these various molecules operate spatiotemporally and how the multiple signaling pathways are coordinated to orchestrate synaptic differentiation are important issues that deserve future investigations. In this line, MAGI-1c, the PDZ-containing scaffolding molecule associated with the C terminus of MuSK is potentially central in the organization of the subsynaptic signaling complex.

Acknowledgments

We thank Pr C. Legay for critical reading of the manuscript, members of the laboratory and the two referees for helpful advice.

References

1. Burden SJ. 2002. Building the vertebrate neuromuscular synapse. *J Neurobiol* 53:501–511.
2. Legay C. 2000. Why so many forms of acetylcholinesterase? *Microsc Res Tech* 49:56–72.
3. McMahan UJ. 1990. The agrin hypothesis. *Cold Spring Harb. Symp Quant Biol* Vol. LV:407–418.
4. Fischbach GD, Rosen KM. 1997. ARIA: a neuromuscular junction neuregulin. *Annu Rev Neurosci* 20:429–458.

5. Sanes J, Litchman JW. 2001. Induction, assembly, maturation and maintenance of a postsynaptic apparatus. *Nat Rev Neurosci* 2:791–805.
6. Ruegg MA, Bixby JL. 1998. Agrin orchestrates synaptic differentiation at the vertebrate neuromuscular junction. *Trends Neurosci* 21:22–27.
7. Meier T, Hauser DM, Chiquet M, Landmann L, Ruegg MA, et al. 1997. Neural agrin induces ectopic postsynaptic specializations in innervated muscle fibers. *J Neurosci* 17:6534–6544.
8. Jennings CG, Dyer SM, Burden SJ. 1993. Muscle-specific trk-related receptor with a kringle domain defines a distinct class of receptor tyrosine kinases. *Proc Natl Acad Sci USA* 90:2895–2899.
9. Valenzuela DM, Stitt TN, DiStefano PS, Rojas E, Mattsson K, et al. 1995. Receptor tyrosine kinase specific for the skeletal muscle lineage: expression in embryonic muscle, at the neuromuscular junction, and after injury. *Neuron* 15:573–584.
10. Bowen DC, Park JS, Bodine S, Stark JL, Valenzuela DM, et al. 1998. Localization and regulation of MuSK at the neuromuscular junction. *Dev Biol* 199:309–319.
11. DeChiara TM, Bowen DC, Valenzuela DM, Simmons MV, Poueymirou WT, et al. 1996. The receptor tyrosine kinase, MuSK, is required for neuromuscular junction formation in vivo. *Cell* 85:501–512.
12. Gautam M, Noakes PG, Moscoso L, Rupp F, Scheller RH, et al. 1996. Defective neuromuscular synaptogenesis in agrin-deficient mutant mice. *Cell* 85:525–535.
13. Jones G, Meier T, Lichtsteiner M, Witzemann V, Sakmann B, et al. 1997. Induction by agrin of ectopic and functional postsynaptic-like membrane in innervated muscle. *Proc Natl Acad Sci USA* 94:2654–2659.
14. Gramolini AO, Burton EA, Tinsley JM, Ferns MJ, Cartaud A, et al. 1998. Muscle and neural isoforms of agrin increase utrophin expression in cultured myotubes via a transcriptional regulatory mechanism. *J Biol Chem* 273:736–743.
15. Lacazette E, Le Calvez S, Gajendran N, Brenner HR. 2003. A novel pathway for MuSK to induce key genes in neuromuscular synapse formation. *J Cell Biol* 161:727–736.
16. Cohen I, Rimer M, Lomo T, McMahan UJ. 1997. Agrin-induced postsynaptic-like apparatus in skeletal muscle fibers in vivo. *Mol Cell Neurosci* 9:237–253.
17. Jones G, Moore C, Hashemolhosseini S, Brenner HR. 1999. Constitutively active MuSK is clustered in the absence of agrin and induces ectopic postsynaptic-like membranes in skeletal muscle fibers. *J Neurosci* 19:3376–3383.
18. Herbst R, Burden SJ. 2000. The juxtamembrane region of MuSK has a critical role in agrin-mediated signaling. *EMBO J* 19:67–77.
19. Lin W, Burgess RW, Dominguez B, Pfaff SL, Sanes JR, et al. 2001. Distinct roles of nerve and muscle in postsynaptic differentiation of the neuromuscular synapse. *Nature* 410:1057–1064.
20. Yang X, Arber S, William C, Li L, Tanabe Y, et al. 2001. Patterning of muscle acetylcholine receptor gene expression in the absence of motor innervation. *Neuron* 30:399–410.
21. Glass DJ, Bowen DC, Stitt TN, Radziejewski C, Bruno J, et al. 1996. Agrin acts via a MuSK receptor complex. *Cell* 85:513–523.
22. Hopf C, Hoch W. 1998. Dimerization of the muscle-specific kinase induces tyrosine phosphorylation of acetylcholine receptors and their aggregation on the surface of myotubes. *J Biol Chem* 273:6467–6473.
23. Glass DJ, Apel ED, Shah S, Bowen DC, DeChiara TM, et al. 1997. Kinase domain of the muscle-specific receptor tyrosine kinase (MuSK) is sufficient for phosphorylation but not clustering of acetylcholine receptors: required role for the MuSK ectodomain? *Proc Natl Acad Sci USA* 94:8848–8853.
24. Zhou H, Glass DJ, Yancopoulos GD, Sanes JR. 1999. Distinct domains of MuSK mediate its abilities to induce and to associate with postsynaptic specializations. *J Cell Biol* 146:1133–1146.
25. Mittaud P, Marangi PA, Herb-Vögltl S, Fuhrer C. 2001. Agrin-induced activation of acetylcholine receptor-bound Src family kinases requires rapsyn and correlates with acetylcholine receptor clustering. *J Biol Chem* 276:14505–14513.
26. Borges LS, Ferns M. 2001. Agrin-induced phosphorylation of the nicotinic acetylcholine receptor regulates cytoskeletal anchoring and clustering. *J Cell Biol* 153:1–12.
27. Smith CL, Mittaud P, Prescott ED, Fuhrer C, Burden SJ. 2001. Src, Fyn, and Yes are not required for neuromuscular synapse formation but are

- necessary for stabilization of agrin-induced clusters of acetylcholine receptors. *J Neurosci* 21:3151–3160.
28. Gautam M, Noakes PG, Mudd J, Nichol M, Chu GC, et al. 1995. Failure of postsynaptic specialization to develop at neuromuscular junctions of rapsyn-deficient mice. *Nature* 377:232–236.
 29. Apel ED, Glass DJ, Moscoso LM, Yancopoulos GD, Sanes JR. 1997. Rapsyn is required for MuSK signaling and recruits synaptic components to a MuSK-containing scaffold. *Neuron* 18:623–635.
 30. Fuhrer C, Hall ZW. 1996. Functional interaction of Src family kinases with the acetylcholine receptor in C2 myotubes. *J Biol Chem* 271:32474–32481.
 31. Qu Z, Apel ED, Doherty CA, Hoffman PW, Merlie JP, et al. 1996. The synapse-associated protein rapsyn regulates tyrosine phosphorylation of proteins colocalized at nicotinic acetylcholine receptor clusters. *Mol Cell Neurosci* 8:171–184.
 32. Dai Z, Luo X, Xie H, Peng HB. 2000. The actin-driven movement and formation of acetylcholine receptors clusters. *J Cell Biol* 150:1321–1334.
 33. Weston C, Yee B, Hod E, Prives J. 2000. Agrin-induced acetylcholine receptor clustering is mediated by the small guanosine triphosphatases Rac and Cdc42. *J Cell Biol* 150:205–212.
 34. Weston C, Gordon C, Teressa G, Hod E, Ren XD, et al. 2003. Cooperative regulation by Rac and Rho of agrin-induced acetylcholine receptor clustering in muscle cells. *J Biol Chem* 278:6450–6455.
 35. Finn AJ, Feng G, Pendergast AM. 2003. Postsynaptic requirement for Abl kinases in assembly of the neuromuscular junction. *Nat Neurosci* 7:717–723.
 36. Pendergast AM. 2002. The Abl family kinases: mechanisms of regulation and signaling. *Ad Cancer Res* 85:51–100.
 37. Luo ZG, Wang Q, Zhou JZ, Wang J, Luo Z, et al. 2002. Regulation of AChR clustering by Dishevelled interacting with MuSK and PAK 1. *Neuron* 35:489–505.
 38. Luo ZG, Je HS, Wang Q, Yang F, Dobbins JC, et al. 2003. Implication of geranylgeranyltransferase I in synapse formation. *Neuron* 40:703–717.
 39. Wang J, Jing Z, Zhang L, Zhou G, Braun J, et al. 2003. Regulation of acetylcholine receptor clustering by the tumor suppressor APC. *Nat Neurosci* 6:1017–1018.
 40. Banks GB, Fuhrer C, Adams ME, Froehner SC. 2003. The postsynaptic submembrane machinery at the neuromuscular junction: requirement for rapsyn and the utrophin/dystrophin-associated complex. *J Neurocytol* 32:709–726.
 41. Cartaud A, Coutant S, Petrucci TC, Cartaud J. 1998. Evidence for in situ and in vitro association between beta-dystroglycan and the subsynaptic 43K rapsyn protein. Consequence for acetylcholine receptor clustering at the synapse. *J Biol Chem* 273:11321–11326.
 42. Grady RM, Akaaboune M, Cohen AL, Maimone MM, Lichtman JW, et al. 2003. Tyrosine-phosphorylated and nonphosphorylated isoforms of alpha-dystrobrevin: roles in skeletal muscle and its neuromuscular and myotendinous junctions. *J Cell Biol* 160:741–752.
 43. Grady RM, Zhou H, Cunningham JM, Henry MD, Campbell KP, et al. 2000. Maturation and maintenance of the neuromuscular synapse: genetic evidence for roles of the dystrophin-glycoprotein complex. *Neuron* 25:279–293.
 44. Stochlic L, Cartaud A, Labas V, Hoch W, Rossier J, et al. 2001. MAGI-1c: A synaptic MAGUK interacting with MuSK at the vertebrate neuromuscular junction. *J Cell Biol* 153:1127–1132.
 45. Dobrosotskaya I, Guy RK, James GL. 1997. MAGI-1, a membrane-associated guanylate kinase with a unique arrangement of protein-protein interaction domains. *J Biol Chem* 272:31589–31597.
 46. Fanning AS, Anderson JM. 1999. Protein modules as organizers of membrane structure. *Curr Opin Cell Biol* 4:432–439.
 47. Sheng M, Pak DT. 1999. Ligand-gated ion channel interactions with cytoskeletal and signaling proteins. *Annu Rev Physiol* 62:755–778.
 48. Schaeffer L, de Kerchove d'Exaerde A, Changeux J-P. 2001. Targeting transcription to the neuromuscular synapse. *Neuron* 31:15–22.
 49. Tanowitz M, Si J, Yu D-H, Feng G-S, Mei L. 1999. Regulation of neuregulin-mediated acetylcholine receptor synthesis by protein tyrosine phosphatase SHP2. *J Neurosci* 19:9426–9435.
 50. Huang YZ, Zang M, Xiong WC, Luo Z, Mei L. 2003. Erbin Suppresses the MAP Kinase Pathway. *J Biol Chem* 278:1108–1114.
 51. Moore C, Leu M, Müller U, Brenner HR. 2001. Induction of multiple signaling loops by MuSK during neuromuscular synapse formation. *Proc Natl Acad Sci USA* 98:14655–14660.
 52. Stochlic L, Cartaud A, Mejat A, Grailhe R, Schaeffer L, et al. 2004. 14-3-3 γ associates with muscle-specific kinase and regulates synaptic gene transcription at vertebrate neuromuscular synapse. *Proc Natl Acad Sci USA* 101:18189–18194.
 53. Van Hemert MJ, de Steensma HY, van Heusden GPH. 2001. 14-3-3 proteins: key regulators of cell division, signalling and apoptosis. *BioEssays* 23:936–946.
 54. Fu H, Xia K, Pallas DC, Cui C, Conroy K, et al. 1994. Interaction of the protein kinase Raf-1 with 14-3-3 proteins. *Science* 266:126–129.
 55. Muslin AJ, Tanner JW, Allen PM, Shaw AS. 1996. Interaction of 14-3-3 with signaling proteins is mediated by the recognition of phosphoserine. *Cell* 84:889–897.
 56. Jacobson C, Cote PD, Rossi SG, Rotundo RL, Carbonetto S. 2001. The dystroglycan complex is necessary for the stabilization of acetylcholine receptor clusters at neuromuscular junctions and formation of the synaptic basement membrane. *J Cell Biol* 152:435–450.
 57. Arikawa-Hirasawa E, Rossi SG, Rotundo RL, Yamada Y. 2002. Absence of acetylcholinesterase at the neuromuscular junction of perlecan-null mice. *Nat Neurosci* 5:119–123.
 58. Kimbell LM, Ohno K, Engel AG, Rotundo RL. 2004. C-terminal and heparin-binding domains of collagenic tail subunit are both essential for anchoring acetylcholinesterase at the synapse. *J Biol Chem* 279:10997–11005.
 59. Cartaud A, Stochlic L, Guerra M, Blanchard B, Lambergeon M, et al. 2004. MuSK is required for anchoring acetylcholinesterase at the neuromuscular junction. *J Cell Biol* 165:505–515.
 60. Engel AG, Ohno K, Sine SM. 2003. Sleuthing molecular targets for neurological diseases at the neuromuscular junction. *Nat Rev Neurosci* 4:339–352.
 61. Liyanage Y, Hoch W, Beeson D, Vincent A. 2002. The agrin/muscle-specific kinase pathway: new targets for autoimmune and genetic disorders at the neuromuscular junction. *Muscle Nerve* 25:4–16.
 62. Chevessier F, Faraut B, Ravel-Chapuis A, Richard P, Gaudon K, et al. 2004. MUSK, a new target for mutations causing congenital myasthenic syndrome. *Hum Mol Genet* 24:3229–3240.
 63. Hoch W, McConville J, Helms S, Newsom-Davis J, Melms A, et al. 2001. Autoantibodies to the receptor tyrosine kinase MuSK in patients with myasthenia gravis without acetylcholine receptors antibodies. *Nat Med* 7:365–368.